



Natural Communities of Louisiana

Calcareous Prairie

Rarity Rank: S1/G1

Synonyms: Barrens, Calcareous Barrens, Calcareous Clay Prairie, Keiffer Prairie, Jackson Prairie, Blackland Prairie, Calcareous Glade

Ecological Systems: CES203.379 West Gulf Coastal Plain Southern Calcareous Prairie



General Description:

- Small, naturally treeless areas ranging in size from less than one acre, up to 80 or more acres
- Occur in a mosaic with calcareous forests
- Occurs on calcareous substrata associated with four geological formations:
 - Fleming Formation (Tertiary-Miocene) in central-western LA
 - Jackson Formation (Tertiary-Eocene) in central LA
 - Cook Mountain Formation (Tertiary-Eocene) in central and western LA
 - Pleistocene Red River terraces in northwest LA (morse clay prairies)
- Soils are stiff calcareous clays (surface pH ~ 7.5-8.0), with very high shrink-swell characteristics, and range in color from red to olive-tan to gray-black
- Various soil inclusions occur (depending on geology) and may include calcareous concretions (limestone nodules), marine mollusc shells, shark teeth, and gypsum crystals
- Regularly-occurring fire, high soil pH, and extreme physical soil properties are believed to generate and perpetuate these upland clay prairies
- Herbaceous flora is very diverse and dominated by grasses, composites, and legumes

Plant Community Associates

Common grass species include:

Schizachyrium scoparium (little bluestem),
Andropogon gerardii (big bluestem),
Aristida spp. (three-awn grasses),
Panicum spp. (panic grasses),
Setaria spp. (bristle grasses)

Sporobolus spp. (dropseeds),
Sorghastrum nutans (Indian grass),
Paspalum spp. (pasp grasses),
Eragrostis spp. (love grasses),

Common composite species include:

Eurybia spp. & *Symphotrichum* spp. (asters),
Liatris spp. (blazing-stars),
Solidago spp. (goldenrods),
Vernonia spp. (ironweeds),
Eupatorium spp. (thoroughworts),
Echinacea purpurea (purple coneflower, rare),
Cacalia plantaginea (Indian plantain),

Helenium spp. (sneeze-weeds),
Coreopsis spp. (tick-seeds),
Ambrosia psilostachya (western ragweed),
Rudbeckia spp. (brown-eyed susans),
Echinacea pallida (pale coneflower),
Silphium spp. (rosin-weeds),
Gaillardia aestivalis (blanket flower)



Natural Communities of Louisiana

Common legumes and forb(wildflower) species include:

Acacia angustissima (prairie acacia),
Desmanthus illinoensis (wad o'pods),
Mimosa strigillosa (sensitive-briar),
Petalostemum candidum (white prairie clover),
Anemone berlandieri (wind flower),
Asclepias spp. (milk-weeds),
Delphinium carolinianum (larkspur),
H. purpurea var. *calycosa* (prairie bluets, rare),
Oenothera speciosa (Mexican evening-primrose),
Salvia azurea (blue sage)

Baptisia spp. (indigos),
Galactia volubilis (milk pea),
Neptunia lutea (yellow puff),
P. purpureum (purple prairie clover),
Ranunculus spp. (crow-foot),
Callirhoe papaver (poppy-mallow),
Hedyotis nigricans (bluets),
Linum spp. (flax),
Ruellia humilis (wild petunia),

Common woody species (invade prairies in absence of fire) include:


Crataegus spp. (hawthorns),
Berchemia scandens (rattan-vine),
Cornus drummondii (rough-leaf dogwood),
Ilex decidua (deciduous holly),
Fraxinus americana (white ash),

Bumelia lanuginosa (chittumwood),
Diospyros virginiana (persimmon),
Juniperus virginiana (eastern red cedar),
Smilax bona-nox (greenbriar),
Gleditsia triacanthos (honeylocust)

Federally-listed plant & animal species: None

Range:

Upper and Lower West Gulf Coastal Plain ecoregions, primarily in central and west-central Louisiana with limited area remaining in northwest LA, and historically known from the Mississippi River Alluvial Plain ecoregion in northeast LA.



Threats & Management Considerations:

Historically there was an estimated 2,000 to 10,000 acres of calcareous prairie statewide but only five to 10 percent of the original extent is thought to remain today. As with the associated calcareous forests, calcareous prairies have mainly been lost to land use changes. Conversion to agriculture or pine plantations and fire suppression represent the greatest loss, while construction of roads, pipelines and utilities, invasive and exotic species, physical damage from timber harvesting, contamination by chemicals (herbicides, fertilizers), and off-road vehicle use all threaten remaining calcareous prairies.

Use of appropriate management activities and developing a compatible management plan prevents destruction or degradation of this habitat type and promotes long-term maintenance of healthy calcareous prairies. Such management strategies should include:

- Use of periodic prescribed fire (every 3 to 5 years)
- Monitoring for and removal of any invasive or exotic species by spot herbicide treatments or mechanical means
- Preventing use of prairie openings as logging sets
- Prohibiting off-road vehicle use or restricting use to existing trails