



Crawfish are freshwater crustaceans found worldwide, except in most tropical regions. In the United States there are 330 known crawfish species, 39 of which can be found in Louisiana. Only two Louisiana crawfish species are targeted for commercial harvest, the red swamp crawfish and the southern white river crawfish.



# Red Swamp Crawfish (Procambarus clarkii)

The red swamp crawfish is the primary commercial species and perhaps the most easily recognized Louisiana crawfish, making up 70-80% of our state's annual catch. Like its name implies, they prefer swampy habitats. Their native range is northeastern Mexico and south-central United States, although they have been introduced to many other countries.



# Southern White River Crawfish (Procambarus zonangulus)

The southern white river crawfish makes up 20-30% of our state's annual crawfish catch. They prefer flooded wetland habitats with periodically flowing, well-oxygenated water. Their native range is Texas, Louisiana, Mississippi, and Alabama, and they only occur in the United States.



Length of day, water temperature, and changes in water level trigger the mating season for Louisiana crawfish species. Females must be slightly soft to mate, usually a few days after molting. Female crawfish mate in open water with males before moving into burrows to spawn. Fertilized eggs are attached under the tail and are carried for two to eight weeks before hatching. Larval crawfish remain on the female for one to three weeks until she leaves the burrow.

## **Habitat**

Crawfish are found in burrows, ditches, rivers, lakes, and streams. Louisiana's commercial crawfish species retreat to burrows when surface waters dry up. These are simple, shallow burrows, less than 40 inches deep. More complex burrows tend to have large chimneys of rounded mudballs capping the entrance, often up to a foot high.

### Diet


Crawfish are opportunistic omnivores that eat almost anything. They can detect small amounts of blood and decomposition chemicals in the water. They feed on partially decomposed plant material, living soft plants, scavenge on any type of animal meat, and are often cannibalistic.

Crawfis

# **Age and Growth**

Crawfish grow by molting, a process in which a new shell forms and causes the outer shell to shed. Molting occurs at night in a protected area away from predators because the new shell is soft, leaving the crawfish vulnerable for several days until the new shell hardens. Juvenile crawfish molt as often as once a week during their first few months, but less often as they mature. Crawfish reach maturity at an age of two months to one year and can live up to three years of age.





### **Red Swamp** Crawfish

- Brown to bright red in color
- Blackish-brown stripe down the abdomen
- More flattened claws



### **Southern White River Crawfish**

- Various shades of brown in color
- No stripe down the abdomen
- More elongated claws



Your purchase of fishing equipment and motor boat fuels supports Sport Fish Restoration and boating access facilities.

> Louisiana Department of Wildlife and Fisheries Office of Fisheries 2000 Quail Drive Baton Rouge, LA 70808 225-765-2800 www.wlf.louisiana.gov

For more information on crawfish: Brian Alford 225-765-2332

This public document was published at a total cost of \$344.60. Five hundred copies of this public document were published and partially paid for by a Federal Aid in Sportfish Restoration Grant (F-136) from the U.S. Fish and Wildlife Service in this first printing at a cost of \$344.60. This document was published by the Louisiana Department of Wildlife and Fisheries, 2000 Quail Drive, Baton Rouge, LA 70808, to inform Louisiana residents and non-residents about the Sport Fish Restoration Program. This material was printed in accordance with standards for printing by state agencies

# **Quick Facts**

There are five species of crawfish that occur only in Louisiana:

- -Teche Painted Crawfish
- -Calcasieu Painted Crawfish
- -Ouachita Fencing Crawfish
- -Caddo Chimney Crawfish
- -Calcasieu Creek Crawfish

Some types of crawfish can burrow over six feet deep.

At least 80% of all crawfish produced in Louisiana are consumed in the state.

Louisiana has more than 1,000 crawfish farmers and over 800 commercial fishermen who catch wild crawfish.

Over 110 million pounds of crawfish are harvested each year in Louisiana, with an economic impact of 120 million dollars.

# Crawfish

### pumps and control water levels.

profitable, requiring less energy to run

**Crawfish Pond Culture** 

accounted for nearly 87% of Louisiana

crawfish production. Crawfish can be

raised in dedicated artificial ponds, but

involves alternating rice with crawfish

In 2010, pond-raised crawfish

generally crawfish pond culture

and sometimes soybeans. This

multiple crop scheme is more

Rice is planted and fields gradually flooded. Crawfish are either already in the field or enter shortly after from neighboring canals and bayous.

March

### March - July

Crawfish are harvested by traps, while the rice grows. Mating also occurs during this time.

#### **August - October**

The rice matures, ponds are drained for harvest, and crawfish retreat to burrows to lay eggs.

#### **October**

The fields are flooded again and crawfish come out from burrows along with new juveniles.

#### **November - January**

Crawfish are harvested. Harvestable crawfish are at least three inches in length.

#### January - March

Cold weather halts feeding and growth.