

## Louisiana Department of Wildlife and Fisheries Natural Heritage Program

## Plethodon websteri Webster's Salamander



**Identification:** A brown salamander with a light red/orange stripe running down its back. Sides are gray or dark brown and the belly is mottled with orange or red patches.

Measurements: Adult: total length 2.75-3 inches. Hatchlings: 13 mm from head to the base of the tail.

**Taxonomic comments:** Although identical in appearance to the zigzag salamander, *Plethodon dorsalis dorsalis*, does not share geographic range and can be separated by molecular analysis.

Status: Global rank is G3 and state rank is S1.

Habitat: Moist hardwood forest bordering rocky streams where they hide beneath rocks, logs, and leaves.

Range: Known from east central Alabama and west central Georgia. Several disjunct populations occur throughout

Alabama, Mississippi, South Carolina, and Louisiana.

Food habits: small arthropods

**Reproduction:** Webster's salamanders reach sexual maturity at around 2 years. Mating begins in January. Pregnant females lay between 3-8 eggs in June and July; eggs hatch between August and September, and hatchlings emerge in October and November.

**Reason for decline:** Habitat destruction including the conversion of hardwood forest to pine plantations.

**Interesting facts:** 1) This salamander is rare in Louisiana, the population covers less than 10 acres. 2) Webster's Salamander is a species of woodland salamander, the most abundant group of invertebrates in forest habitats. These species are important indicators of habitat quality and degradation because of their concentrated populations, dependence on both terrestrial and aquatic habitats, and role as insect predators. Declines in salamander populations are early indications of ecosystem damage that may still be reversed.

## **Known distribution in Louisiana:**



Dates of documented observations are: 1999

## References

- Camp, C.D., and L.L. Bozeman. 1981. Foods of two species of <u>Plethodon</u> (<u>Caudata: Plethodontidae</u>) from Georgia and Alabama. Brimleyana 6:163-166.
- Conant, R. and J. T. Collins. 1991. A Field Guide to Reptiles and Amphibians: Eastern and Central North America. Houghton Mifflin Company, Boston, Massachusetts.
- Dundee, H. A., and D. A. Rossman. 1989. The Amphibians and Reptiles of Louisiana. Louisiana State University Press, Baton Rouge, Louisiana.
- Herbeck, L.A. and R.D. Semlitsch. 2000. Life history and ecology of the Southern Redback Salamander, *Plethodon serratus*, in Missouri. Journal of Herpetology 34:341-347.
- Larson, D. L. 1998. Tiger salamander life history in relation to agriculture in the northern Great Plains: A hypothesis. *In* Status and conservation of Midwestern amphibians. Lannoo, E. J. ed. University of Iowa Press, Iowa City, Iowa.
- NatureServe Explorer: An online encyclopedia of life [web application]. 2002. Version 1.6. Arlington, Virginia, USA: NatureServe. Available: http://www.natureserve.org/explorer. (Accessed: January 2, 2003).
- Semlitsch, R.D. and C.A. West. 1983. Aspects of the life history and ecology of Webster's Salamander, *Plethodon websteri*. Copeia 2:339-346.
- Wilson, L. A. 1995. Land Manager's Guide to the Amphibians and Reptiles of the South. Nature Conservancy, Chapel Hill, North Carolina